


AMERICAN ASSOCIATION OF SUICIDOLOGY

Youth Suicide Fact Sheet Based on 2011 Data (2013)

Overview

- In 2011, 4, 822 youth age 15-24 died by suicide.1
- Suicide was the 2nd leading cause of death for 15 - 24 year-olds in 2011
- The 2011 Youth Risk and Behavior Survey found that in the previous 12 months among high school students; 15.8% seriously considered suicide; 12.8% made a plan for suicide; 7.8% attempted suicide one or more times; 2.4% made a suicide attempt that had to be treated by doctor or nurse.

Suicide Rate by Race,



Girls are more likely to attempt suicide, but boys are 4.34 times more likely to die by suicide than girls.

Risk Factors

- Mental Illness
- Substance abuseiii
- Firearms in the household^v
- Previous suicide attemptsvii
- Non-suicidal self-injuryix
- Exposure to friends'/family members'
- Low self-esteemxii

Protective Factors

- Family and school connectednessⁱⁱ
- Safe schoolsiv
- Reduced access to firearmsvi
- Academic Achievementviii
- Self-esteem^x

Youth Suicide Can Be Prevented

Youth across America are affected by suicide, be it in families, at school, or in their own lives. The cost is immesasurable, but help is available.

Friends are often the first to know their friend is suicidal, and we need to help them know where to find help.

Source: 'Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based Injury Statistics Query and Reporting System (WISQARS) [online]. [cited February 2012]. Available from www.cdc.gov/ncipc/wisqars)

ii Kaminski et al, J Youth Adol, 2010 iii Brent et al., J Am Acad Child Adol Psych, 1999. iv Eisenberg et al., J PED, 2007

^vBrent et al., J Am Acad Child Adol Psych, 1999

vi Grossman et al., JAMA, 2005

vii Beautrais, SLTB, 2004

viii Borowsky et al., Pediatrics, 2001

ix Nock et al., Psychi Res, 2006

x Sharaf e al., JCAPN, 2009

xi Borowsky et al., Pediatrics, 2001

xii Resnick et al., JAMA, 1997